

EVALUATION OF ALPHA-AMYLASE INHIBITORY ACTIVITY OF *GREWIA ABUTILIFOLIA* LEAVES**Abdullah-Al-Mamun*, Md. Shariful Islam, Md. Ibrahim Sunny, Md Abdur Rahaman Miah, Umma Hafsa, Shakila Akter Zabin, Pankoj Chandra Roy**

Department of Pharmacy, Southeast University, Banani, Dhaka-1213, Bangladesh.

***Corresponding Author: Abdullah-Al-Mamun**

Department of Pharmacy, Southeast University, Banani, Dhaka-1213, Bangladesh.

Article Received on 28/08/2017

Article Revised on 19/09/2017

Article Accepted on 10/10/2017

ABSTRACT

Alpha-amylase inhibitors are oral anti-diabetic drugs used for diabetes mellitus type 2 that work by preventing the digestion of carbohydrates (such as starch and table sugar). Inhibition of Alpha-amylase enzyme that plays a role in digestion of starch and glycogen is considered a strategy for the treatment of disorders in carbohydrate uptake such as diabetes and obesity. Plants are an important source of chemical constituents with potential for inhibition of Alpha-amylase and can be used as therapeutic or functional food sources. *Grewia abutilifolia* belonging to the family Tiliaceae, is a tree and that is used in traditional medicine for the treatment of diabetic mellitus. In the present study, methanolic extracts of *Grewia abutilifolia* leaves which are used in the Ayurvedic traditional system of medicine to treat diabetes were tested for their inhibitory effect on α -amylase. The crude methanol extracts (6.88 ± 0.14) of *Grewia abutilifolia* with pet-ether fraction, (21.13 ± 25.24), chloroform fraction (6.02 ± 1.15) and methanol+water fraction (5.2 ± 0.80). Among all fractions acarbose & methanol+water showed highest percentage inhibition & the IC_{50} value was 5.7 ± 0.24 & 5.2 ± 0.80 . So, methanol + water extracts has the highest percentage of inhibition than acarbose. As a results, *Grewia abutilifolia* leaves properly inhibits the alpha-amylase in diabetes.

KEYWORDS: Alpha-amylase, Carbohydrate Metabolism, Diabetes Mellitus, *Grewia abutilifolia*.**INTRODUCTION**

Now a days, the dependency of the medicine on plants materials is becoming clearer. However, this dependency upon medicinal plants is still not widely acknowledged.^[1] Plants possess a large number of vital compounds that might form a part of healthy diet. Traditionally, different parts of plants were used directly as pharmaceuticals.^[2] Clinically useful chemicals are now being obtained from plants even those that have not been classified before as medicinal plants. One of the strategies and methods adopted to cure diabetes mellitus involves the inhibition of carbohydrate digesting enzymes such as amylase.^[3-4] Alpha-amylase inhibitors are oral anti-diabetic drugs used for diabetes mellitus type 2 that work by preventing the digestion of carbohydrates (such as starch and table sugar).^[5] Carbohydrates are normally converted into simple sugars (monosaccharides) which can be absorbed through the intestine.^[6] Hence, alpha-glucosidase inhibitors reduce the impact of carbohydrates on blood sugar.^[7] They are used to establish greater glycemic control over hyperglycemia in diabetes mellitus type 2, particularly with regard to postprandial hyperglycemia.^[8] They may be used as monotherapy in conjunction with an appropriate diabetic diet and exercise or they may be used in conjunction with other anti-diabetic drugs.^[9] *Grewia Abutilifolia* plant leaves are found in Bangladesh.^[10-13] It is used as a remedy for acne and a

root decoction as a remedy for fever.^[14] The bark yields good quality fibre that is made into rope.

MATERIALS AND METHODS**Collection, Identification & authentication of plant**

The plant material (leaves) was collected during the month of March, 2015 from Comilla Hill track, Bangladesh in fresh condition and identified by an expert taxonomist. A voucher specimen was submitted to the National Herbarium Bangladesh (accession number: 41884). Leaves were then washed properly to remove dirty materials and shade dried for several days with occasional sun drying. These were then dried in an oven for 24 hours at considerably low temperature for better grinding. The dried leaves were ground into coarse powder by a grinding machine in the department of Pharmacy, Southeast University.

Extraction of the plant materials: Powdered plant materials (leaves) having a weight of about 500 gm. were taken in an amber colored reagent bottle and soaked in 1.5 liter of methanol. The bottle with its contents were sealed and kept for a period of about 7 days with occasional shaking and stirring. The whole mixture was then filtered through cotton and then through Whatman No.1 filters paper and was concentrated with a rotary

evaporator under reduced pressure at 52°C temperature to afford crude extract (45.39 gm).

Solvent-solvent partitioning of crude extract (*Grewia Abutilifolia*): An aliquot (20gm) of the concentrated methanol extract was fractionated by modified Kupchan method with petroleum ether, chloroform and water and the amount of fractions obtained is given in the following table:

Table. 1: Different fractions with amount of *Grewia abutilifolia*.

Name of fractions	Weight of fractions (gm)
Petroleum ether fraction	7.37
Chloroform fraction	7.25
Aqueous fraction	5.38

Alpha-amylase inhibition test

Principle

^[15]Diabetes mellitus is a complex disease that is characterized by gross derangement in carbohydrate, protein, and fat metabolism.^[16]It is a progressive metabolic disorder of glucose metabolism that eventually leads to micro and macro vascular changes causing secondary complications that are difficult to manage.^[17]Type-I diabetes results from inadequate synthesis of insulin by beta -cells of the pancreas while type II diabetes is characterized primarily by insulin resistance (a condition in which peripheral cells do not respond normally to insulin) or beta-cell dysfunction.^[18]Alpha-amylase is a prominent enzyme found in the pancreatic juice and saliva which breaks down large insoluble starch molecules into absorbable molecules.^[19]On the other hand, mammalian alpha-glucosidase in the mucosal brush border of the small intestine catalyzes the end step of digestion of starch and disaccharides that are abundant in human diet.^[20]

Inhibitors of alpha-amylase and alpha-glucosidase delay the breaking down of carbohydrates in the small intestine and diminish the postprandial blood glucose excursion.

Preparation of 0.5M Tris-Hcl buffer (pH 6.9)

^[21-22]Tris – Hcl was added with calcium chloride in 800 ml of water. Then added slowly 200ml of water. Then check the pH by the pH meter. If the pH was basic added acid & if the pH was acidic added base.

Preparation of starch solution

100mg starch was mixed with 10ml Tris Hcl buffer.

Preparation of α -amylase solution: 0.007gm α -amylase was dissolved in 100ml of Tris Hcl buffer.

Preparation of 50% acetic acid solution: 50gm / 50ml acetic acid was dissolved in 100ml of water.

To conduct this experiment, 8 test tube were taken.^[23] 0.6 ml starch solution taken in each test tube Boiled at 90°C for 10 minute. Then pre incubate at 37°C for 5 minute. 0.6ml of sample/standard solution was added to each test tube.^[24] 0.3ml of α -amylase solution is incubate at 37°C for 10 minutes. Then added 1.5ml of 50% acetic acid solution. Reaction mixture was centrifuged at 3000 rpm for 8 minutes. Then taken absorbance of supernatant at 595 nm.

RESULTS AND DISCUSSION

Alpha-amylase inhibition studies demonstrated that the extracts of *Grewia abutilifolia* leaf had inhibitory activity. In all experiments, control samples were prepared accordingly without any plant extract and were compared with the test samples containing various concentrations of the plant extract. Acarbose was used as the standard.

Table. 02: Alpha-amylase inhibitor activity of different fractions of *Grewia Abutilifolia* at different concentrations.

Name of sample	Conc. ($\mu\text{g/ml}$)	% of inhibition			% of inhb. Mean	% of inhb. \pm STD	IC ₅₀ ($\mu\text{g/ml}$)
		A	b	c			
CME	7.81	58.07	56.25	55.87	56.73	1.18	6.88 \pm 0.14
	15.62	60.05	72.13	60.36	64.18	6.89	
	31.25	64.69	74.36	67.14	68.73	5.03	
	62.5	69.69	80.45	74.25	74.80	5.40	
	125	79.04	86.15	79.25	81.48	4.05	
	250	86.12	90.25	82.19	86.19	4.03	
Pt-Ether fraction	7.81	73.37	50.24	49.85	57.82	13.47	21.13 \pm 25.24
	15.62	80.17	55.24	53.65	63.02	14.87	
	31.25	86.66	64.49	62.25	71.13	13.49	
	62.5	86.97	70.25	70.25	75.82	9.65	
	125	86.12	72.19	82.29	80.20	7.20	
	250	92.92	76.37	86.19	85.16	8.32	
Chloroform fraction	7.81	70.25	53.25	75.24	66.25	11.53	6.02 \pm 1.15
	15.62	75.36	60.15	62.03	65.85	8.29	
	31.25	79.05	64.25	75.45	72.92	7.72	
	62.5	82.72	79.45	82.15	81.44	1.75	
	125	83.85	83.25	88.05	85.05	2.62	

	250	85.55	88.19	90.25	88.00	2.36	
Methanol + water fraction	7.81	90.93	70.25	67.45	76.21	12.82	5.2 ± 0.80
	15.62	91.76	74.02	70.25	78.68	11.49	
	31.25	92.06	81.5	76.15	83.24	8.10	
	62.5	92.35	89.15	89.25	90.25	1.82	
	125	93.48	90.25	86.12	89.95	3.69	
	250	95.75	94.48	87.25	92.49	4.59	

Table. 3: Alpha-amylase inhibitor activity of Acarbose(Standard)at different concentrations.

Name of sample	Conc. (µg/ml)	% of inhibition			% of inh. Mean	% of inh. ± STD	IC ₅₀ (µg/ml)
		a	b	c			
Acarbose (Standard)	7.81	70.27	65.27	69.87	68.47	2.78	5.7 ± 0.24
	15.62	71.15	66.15	61.45	66.25	4.85	
	31.25	81.32	76.32	70.32	75.99	5.51	
	62.5	83.67	78.67	75.67	79.34	4.04	
	125	93.34	88.34	83.25	88.31	5.05	
	250	95.57	90.57	85.37	90.50	5.10	




Figure. 1: Alpha-amylase inhibitor activity of CME, PEF, CLF, MWF & Acarbose(STD) of *Grewia abutilifolia* at different concentrations.




Figure. 2: IC₅₀ (µg/ml) values of different extractives of *Grewia abutilifolia* for Alpha- amylase activity Assay.

DISCUSSION

[25] Alpha-amylase inhibitors are oral anti-diabetic drugs used for diabetes mellitus type 2 that work by preventing the digestion of carbohydrates (such as starch and table sugar). Carbohydrates are normally converted into simple sugars (monosaccharides) which can be absorbed

through the intestine.^[26-27] Acarbose is used as standard in this result which blocks pancreatic alpha-amylase in addition to inhibiting membrane-bound alpha-glucosidases.^[28] Pancreatic alpha-amylase hydrolyzes complex starches to oligosaccharides in the lumen of the small intestine.^[29] Inhibition of these enzyme systems reduces the rate of digestion of carbohydrates.^[30] Less glucose is absorbed because the carbohydrates are not broken down into glucose molecules.^[31-32] In diabetic patients, the short-term effect of these drugs therapies is to decrease current blood glucose levels, the long-term effect is a small reduction in hemoglobin_{A1c} level.^[33] In alpha amylase inhibition test the crude methanolic extract of *Grewia abutilifolia* with known antidiabetic activity was investigated for their potential to inhibit α -amylase activity.^[34] Among the different fractions of the crude extract such as petroleum ether, chloroform and aqueous fractions, were examined at a different concentration (µg/ml).^[35-36] Among all fractions acarbose & methanol + water showed highest percentage inhibition & the IC₅₀ value was 5.7 ± 0.24 & 5.2 ± 0.80.^[37] So, methanol + water has the highest percentage of inhibition than acarbose. pet-ether showed lowest percentage of inhibition & the IC₅₀ value was 21.13 ± 25.24. Chloroform fraction & Crude Methanol extract showed moderate percentage of inhibition & the IC₅₀ value was 6.02 ± 1.15 & 6.88 ± 0.14 respectively.^[38] So, the crude methanol extract of *Grewia abutilifolia* showed alpha amylase inhibition activity.

CONCLUSION

Diabetes mellitus is a metabolic disorder characterized by chronic hyperglycemia and its type II is the major form of diabetes, accounting for 90% of cases worldwide. There are many and diverse therapeutic strategies in the management of Type II diabetes. The inhibition of carbohydrate hydrolyzing enzymes such as α -amylase can be an important strategy to lower postprandial blood glucose levels. The leaves and bark have been also used in medicated enema and used in traditional medicine for the treatment of diabetic mellitus

and is widely distributed all over the Bangladesh. In this research, *Grewia abutilifolia* are used for alpha amylase inhibition test. The result demonstrates that, In alpha amylase inhibition test among all fractions of *Grewia abutilifolia* acarbose & methanol + water showed highest inhibition respectively. Further studies needed to determine active compounds responsible for these activities and I think that, this project work will help to identify the active constituents, structure and their action mechanism responsible for the activity and this project will also help a new investigator to proceed the future research.

ACKNOWLEDGEMENTS

After all, the authors highly acknowledged the financial support from the Department of Pharmacy, Southeast University, Banani, Dhaka-1213, Bangladesh. Also, We are all thanking to Professor & Chairman Dr. Md. Siddiqui Islam for his valuable directions to carry out our research project.

REFERENCES

- Bhosale U. P, Hallale B. V, Asian Journal of Plant Science and Research, 2011; 1: 96.
- Source:Herbal medicine|University of Maryland Medical Center
<http://umm.edu/health/medical/altmed/treatment/herbal-medicine#ixzz3t3to0XeL>
University of Maryland Medical Center.
- Jump up^ Verrotti A, Scaparrotta A, Olivieri C, Chiarelli F. "Seizures and type 1 diabetes mellitus: current state of knowledge". European journal of endocrinology, December 2012; 167(6): 749–58. doi:10.1530/EJE-12-0699. PMID 22956556.
- Bhise.B,Bhutkar S.M.A. In vitro assay of alpha amylase inhibitory activity of some indigenous plants, 2012; 10(1): 457-46.
- Sharma, H. M.; Bodeker, Gerard C (1997). "Alternative Medicine (medical system)". Encyclopædia Britannica (2008 ed. ed.)
- Falcão HS, Moura MD, Diniz MFFM, Modesto-Filho J. Plants and their active constituents from South, Central, and North America with hypoglycemic activity. Rev Bras Farmacogn, 2005; 15: 392-413.
- Pignone M, Alberts MJ, Colwell JA, Cushman M, Inzucchi SE, Mukherjee D, Rosenson RS, Williams CD, Wilson PW, Kirkman MS. "Aspirin for primary prevention of cardiovascular events in people with diabetes: a position statement of the American Diabetes Association, a scientific statement of the American Heart Association, and an expert consensus document of the American College of Cardiology Foundation". Diabetes Care, June 2010; 33(6): 1395–402. doi:10.2337/dc10-0555. PMC 2875463.PMID 20508233.
- Ripsin CM, Kang H, Urban RJ. "Management of blood glucose in type 2 diabetes mellitus" (PDF). American family physician, 2009; 79(1): 29-36. PMID 19145963.
- M.V.Shrimanker^{1*}, D.D.Patel², H.K.Modi¹, K.P.Patel¹, R.M.Dave.In-vitro Antibacterial and Antifungal Activity of *Grewia Asiatica* Linn. Leaves Saraswati Institute of Pharmaceutical Sciences, Dhanap, Gandhinagar - 382355, Gujarat, India Shri B. M. Shah Pharmaceutical Education and Research, Modasa - 383315, Gujarat, India.
- Penso, G 1980. WHO Inventory of Medicinal Plants Used In Different Countries. Geneva. Switzerland, WHO.
- GOYAL. K. PDepartment of Chemistry, University of Rajasthan, Jaipur, India. PHYTOCHEMICAL AND PHARMACOLOGICAL PROPERTIES OF THE GENUS GREWIA: A REVIEW:30 July 2012.
- Khadeer Ahamed MB¹, Krishna V, Dandin CJ. In vitro antioxidant and in vivo prophylactic effects of two gamma-lactones isolated from *Grewia tiliaefolia* against hepatotoxicity in carbon tetrachloride intoxicated rats.
- Rahmanullah Siddiqi^{1*}, Shahina Naz¹, Samia Ahmad² and Syed Asad Sayeed¹, Antimicrobial activity of the polyphenolic fractions derived from *Grewia asiatica*, *Eugenia jambolana* and *Carissa carandas*. 12 JAN 2011.
- <http://tropical.theferns.info/viewtropical.php?id=Grewia+abutilifolia>.
- Klein.O, Lyngø J, Endahl.L, Damholt.B, Nosek B, and Heise T, "Albumin-bound basal insulin analogues (insulin detemir and NN344): comparable time-action profiles but less variability than insulin glargine in type 2 diabetes," Diabetes, Obesity and Metabolism, 2007; 9(3): 290–299.
- Hyperglycemia (High blood glucose). American Diabetes Association.
<http://www.diabetes.org/living-with-diabetes/treatment-and-care/blood-glucose-control/hyperglycemia.html>. Accessed April 29, 2014.
- Heise T, Nosek .L, Rønn .B et al., "Lower within-subject variability of insulin detemir in comparison to NPH insulin and insulin glargine in people with type 1 diabetes," Diabetes, 2004; 53(6): 1614–1620.
- Cooke DW, Plotnick L. "Type 1 diabetes mellitus in pediatrics". Pediatr Rev., November 2008; 29(11): 374–84., quiz 385. doi:10.1542/pir.29-11-374. PMID 18977856.
- Afifi. A. F., Kamel E. M, Khalil A. A., Foad M. A., Fawziand E. M., and Houseny M., "Purification and characterization of α -amylase from *penicillium olsonii* under the effect of some antioxidant vitamins," Global Journal of Biotechnology and Biochemistry, 2008; 3(1): 14–12.
- Manoha. V N., Talpur, B., Echard W, Lieberman S, and Preuss H. G., "Effects of a water-soluble extract of maitake mushroom on circulating glucose/insulin concentrations in KK mice," Diabetes, Obesity and Metabolism, 2002; 4(1): 43-48.

21. Kwon Y.-I, Apostolidis E., and Shetty K, "Evaluation of pepper (*Capsicum annuum*) for management of diabetes and hypertension," *Journal of Food Biochemistry*, 31(3).
22. Funke I, Melzig MF. Effect of different phenolic compounds on alpha-amylase activity: screening by microplate-reader based kinetic assay. *Pharmazie*, 2005; 60: 796-797.
23. <https://www.hindawi.com/journals/tswj/2017/3592491/>
24. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2959209/>
25. Diabetes mellitus (DM). The Merck Manual for Health Care Professionals. http://www.merckmanuals.com/professional/endocrine_and_metabolic_disorders/diabetes_mellitus_and_disorders_of_carbohydrate_metabolism/diabetes_mellitus_dm.html. Accessed April 29, 2014.
26. Atkinson MA, et al. Type 1 diabetes. *The Lancet*. 2014;383:69.
27. Jump up^ Verrotti A, Scaparrotta A, Olivieri C, Chiarelli F. "Seizures and type 1 diabetes mellitus: current state of knowledge". *European journal of endocrinology*, December 2012; 167(6): 749–58. doi:10.1530/EJE-12-0699. PMID 22956556.
28. Kitabchi AE, Umpierrez GE, Miles JM, Fisher JN. "Hyperglycemic crises in adult patients with diabetes". *Diabetes Care*, July 2009; 32(7): 1335–43. doi:10.2337/dc09-9032. PMC 2699725. PMID 19564476.
29. Sarwar N, Gao P, Seshasai SR, Gobin R, Kaptoge S, Di Angelantonio E, Ingelsson E, Lawlor DA, Selvin E, Stampfer M, Stehouwer CD, Lewington S, Pennells L, Thompson A, Sattar N, White IR, Ray KK, Danesh J (2010). "Diabetes mellitus, fasting blood glucose concentration, and risk of vascular disease: A collaborative meta-analysis of 102 prospective studies"
30. Diabetes mellitus (DM). The Merck Manuals: The Merck Manual for Health Care Professionals. http://www.merckmanuals.com/professional/endocrine_and_metabolic_disorders/diabetes_mellitus_and_disorders_of_carbohydrate_metabolism/diabetes_mellitus_dm.html. Accessed April 29, 2014.
31. Cavanagh PR. "Therapeutic footwear for people with diabetes". *Diabetes Metab. Res. Rev.*, 2004; 20(Suppl 1): S51-5. doi:10.1002/dmrr.435. PMID 15150815.
32. Bhise.B, Bhutkar S.M.A. In vitro assay of alpha amylase inhibitory activity of some indigenous plants, 2012; 10(1): 457-46.
33. Book of the plants extraction procedure and solvent partition method, 2001; 9: 122-248.
34. https://employees.csbsju.edu/cschaller/Principles%20Chem/purification/solvent_partitioning.htm.
35. Nkunya MH¹, Weenen H, Bray DH, Mgani QA, Mwasumbi LB. Antimalarial activity of Tanzanian plants and their active constituents: the genus *Uvaria*. 1991 Aug; 57(4).
36. MEADE. C and PARNELL.J Multivariate analysis of leaf shape patterns in Asian species of the *Uvaria* group (Annonaceae). 10 NOV. 2003.
37. Nkunya MH¹, Weenen H, Bray DH, Mgani QA, Mwasumbi LB. Antimalarial activity of Tanzanian plants and their active constituents: the genus *Uvaria*. 1991 Aug; 57(4).
38. www.theplantlist.org/tpl/record.