

UTILIZATION OF CASSIA SOPHERA L. SEED OIL CAKE AS BIOFERTILIZER.***Dr. Shahin Aziz**

Chemical Research Division, Bangladesh Council of Scientific and Industrial Research (BCSIR) Laboratories, Dhaka, Bangladesh.

***Corresponding Author: Dr. Shahin Aziz**

Chemical Research Division, Bangladesh Council of Scientific and Industrial Research (BCSIR) Laboratories, Dhaka, Bangladesh.

Article Received on 16/04/2018

Article Revised on 06/05/2018

Article Accepted on 26/05/2018

ABSTRACT

Cassia Sophera Linn is one of the medicinally important plants belonging to the family of Leguminosae. The whole part of the plant used as traditional folk medicine for the treatment of pityriasis, psoriasis, asthma, acute bronchitis, cough, diabetes and convulsions of children. anti diathorreal and antipyretic effects. Oil extracted from the seeds of this plant is used as energy source as well as in cosmetics while the cake (a byproduct after extracting oil) was found to be rich in minerals in the form of plant nutrients. In order to reduce the production cost by minimizing or utterly eliminating the use of chemical fertilizers, decreasing environmental hazards, improve soil structure, promote leveraging agriculture and obtain high quality crops, *C. Sophera* seed oil cake getting importance due its rich mineral contents. *Cassia Sophera* seed oil cake was assessed and it was found to be a potential source of bio-fertilizer due to richness of nutrients. The analysis of valuable elements present in this oil cake was done by Atomic absorption spectrophotometer.

KEYWORDS: *Cassia Sophera* seed oil cake, Bio-fertilizer, Atomic Absorption Spectrophotometer, elemental compositions.

1. INTRODUCTION

Cassia Sophera Linn is commonly known as "Kulkasunda" which belongs to the family of Leguminosae. It is one of the important medicinal plants in tropical and subtropical region in Asia especially in India, Sri Lanka, Pakistan, Malaysia, Myanmar and in Bangladesh. It grows abundantly in the plain land, hilly areas of Chittagong Hill Tracts, Sylhet and patches throughout Bangladesh.^[1] The seeds of this important medicinal plant are used as a traditional medicine in Japan, Korea and China for the treatment of eye in Ammation, phytophobia and lacrimation, dysentery, headache as well as dizziness.^[2] The *Cassia Sophera* L seed oil is used as to promote a healthy immune function. It is also a great oil to diffuse during cold months due to its warming properties and spicy scent.^[3] The plant *Cassia Sophera* L revealed the presence of ascorbic acid, dehydroascorbic acid, β -sitosterol, glycosides and a rich source of flavanoids and anthraquinones.^[4] *Cassia Sophera* seed oil cake (a byproduct after extracting oil) is obtained from the seeds of *Cassia Sophera* plant.

The *Cassia Sophera* seed oil cake is a byproduct obtained from *Cassia Sophera* seeds with hot pressing and can be used in the production of biofertilizer due to its richness in mineral contents, which is necessary for plant growth. Although the importance of chemical fertilizers, many constraints have been raised such as their adverse impacts on the public health, environment, increasing the

production cost and deterioration of soil fertility.^[5] It is very important to find alternative methods for supplying nutrients to the growing plants to confront the previous problems. Now a days, various researchers consider the utilization of organic and bio-fertilizers as promising alternative nutrition especially for developing countries.

Organic fertilization provides the means for stabilizing soil fertility (especially in newly reclaimed soils) converting nitrogen in less soluble form is the main advantage of organic fertilizers compared to chemical fertilizers. The favorable effect of cattle manure on the vegetative growth of some medicinal plant has been reported by numerous investigators. The vegetative growth, fruit as well as oil yields were enhanced by using organic manure.^[6,7] Moreover, the photosynthetic pigments and carbohydrate percentage and N, P and K contents^[8,6] were also enhanced. Recent awareness has been offered to reduce pollution practices in sustainable agriculture one of the way to minimize soil pollution is using bio-stimulants compounds without causing any harmful effects on aerial and soil environment to retard nitrification for sufficiently longer time and increase the soil fertility.^[9] The effective utilization of bio-fertilizers for crops not only provides economic benefits to the procedures, but also improves, maintains the soil fertility and sustainability in natural soil eco system. The beneficial effects of plant growth promoting rhizobacteria on growth are not only through Nitrogen

fixed in the rhizosphere, but also related to the ability of these bacteria to synthesize antibiotics and growth-promoting substances including phytohormones and sometimes the ability to solubilize phosphates; the use of phosphates solubilizing bacteria becomes necessary to minimize the dose of chemical P fertilizer that cause environmental pollution.^[10,11]

The mix treatment of nitrogen fixing bacteria and phosphate solubilizing bacteria resulted in the maximum increase in most of the growth and yield parameters of several medicinal and aromatic plants.^[12,13,14,15,11,7] The essential oil content and its main components of various Apiaceae fruits are increased by using bio-fertilizers. In addition, carbohydrate percentage as well as nitrogen, phosphorus and potassium content in leaves were also promoted.^[7,13,16]

The combination of organic manure and bio-fertilizers prove to be the superior treatment compared to the individual application, which reflected in the greatest influence on growth, yield, oil content and chemical constituents of different medicinal and aromatic plants.^[10,17,19]

Although bio-fertilizers are known to play an important role in plant production, little is known about the interactive effects of both of them on growth and secondary metabolites accumulation in medicinal plants. To the best of our knowledge, no systemic studies have been conducted to analyze the mineral contents of *C. Sophera* seed oil cake so far. So, the aim of this study was to investigate the elemental compositions in *Cassia Sophera* seed oil cake and justify the usability of it as a bio-fertilizer.

2. MATERIALS AND METHODS

2.1 Collection of plant material

Fully matured seeds of *Cassia Sophera* were collected from Sylhet, Bangladesh in the month of June 2015 and identified by the taxonomist of Bangladesh national Herbarium, Dhaka where a voucher specimen (No.43734) has been deposited. Then they were cleaned, air dried and seeds were powdered using 20 mesh screen in Willey mill and then used for collecting oil cake after hot extraction of oil from the seeds.

2.2. Reagents and Standards

All reagents used were from Merck (Darmstadt, Germany) or Sigma Aldrich (Buchs, Switzerland). Petroleum ether (b.p 40-60 °C, Merck, Germany) of AR grade, under normal atmospheric pressure was employed for extraction of plant material. Solvent from extract were recovered under distillation and the dried extracts were preserved in a refrigerator.

2.3. Extraction & preparation of *C. Sophera* seed oil cake

The *Cassia Sophera* seeds were cleaned to separate from dirt and grounded to obtain powder so that maximum

particle exposure was got for extraction of oil. About 100 g of powder were extracted with ANALAR petroleum ether (b.p 40°C -60°C) in a soxhlet apparatus for 72h. The extract was first filtered and then vacuum distilled to remove solvent completely. The color of the extracted oil was chocolate maroon. Then the extracted oil was filtered through ANALAR activated charcoal. Finally, *C. Sophera* seed oil cake (a byproduct after getting oil) was obtained. The yield of *Cassia Sophera* seed oil cake was calculated (85%) and stored.

2.4. The proximate Analysis

Cassia Sophera seed oil cake was subjected to determine the percentage of moisture, ash, water soluble and insoluble ash, carbohydrate content according to the standard methods.

2.4.1. Moisture and Dry matter contents

The Moisture content was determined by heating the samples in an electric oven at 105-110°C until constant weight (6-10 hours).^[23] The percentage was calculated by,

$$\text{Moisture content}(\%) = \frac{\text{Weight of moisture}}{\text{Weight of sample taken}} \times 100$$

Dry matter (%) = 100-moisture%

2.4.2. Ash Contents

Ash was determined by incineration of the moisture free samples at about 600°C (about 6-12 hours) in a temperature controlled Muffle furnace until ash becomes almost white or grayish white in color.^[23] The percentage of ash was calculated by,

$$\text{Ash content on dry weight basis}(\%) = \frac{\text{Weight of ash}}{\text{Weight of sample taken}} \times 100$$

2.4.3. Water Soluble Ash

Water soluble ash was determined by boiling the ash sample of *C. Sophera* oil seed cake with 25ml distilled water for 5 minutes and then insoluble matter was thus collected.^[24] Then they were dried, ignited at 450°C and weighed. The percentage from the ash taken was calculated by,

$$\text{Water soluble ash}\% = \frac{(\text{Weight of ash taken} - \text{Weight of water in soluble ash}) \times \text{Ash Content}(\%)}{\text{Weight of ash taken}}$$

2.4.4. Mineral Compositions

Nitric acid (69%, Merck India) and Perchloric acid (70%, Merck India) was used for the digestion of samples to quantify mineral compositions in dried *C. Sophera* seed oil cake. For standard calibration of respective elements Na, K, Ca, Mg, Fe, Zn, Ni, Mn, Cu, Pb standard solution (100mg/ml) from were purchased from Hach (Germany). The respective desired standard from the stock solution using lab made double distilled water were prepared.

2.4.5. Ashing and digestion of plant parts

Accurately 2.0g of *C. Sophera* seed oil cake was taken in a porcelain crucible and heated to about 650°C and cooled and was weighed. The crucible with sample was placed

in the Bunsen burner (at low flow rate gas) until the smoke ceased. Then the crucible was placed in a muffle furnace at 525°C for about 8-10 hours to obtain carbon free white ash. It was then cooled in desiccators and weighed. This procedure was repeated till the color of the ash was changed to almost white as well as constant weight was obtained. About 1.0g ash sample for *C. Sophera* seed oil cake was taken separately in 50ml volumetric flask and then 15ml 1M HNO₃ acid was added. Then the flask was placed on magnetic stirrer heater in fume hood for four hours at 250°C. When the color of the solution was changed to milky solutions, it was cooled for 10 minutes and then 7.5ml concentrated perchloric acid (HClO₄) was added. Then it was heated until colorless solution was obtained. For the determination of dissolved elements, the sample was filtered through 0.45 micron filter paper. In all the cases, the pH of the sample was maintained and verified to be less than 2.0 prior to analysis.^[25] The standard working solution of interest was prepared to make the standard calibration curve. Absorption for a sample solution used the calibration curve to determine the concentration of particular element in that sample.

2.4.6. Analytical procedure

Among all elements only Sodium (Na) and Potassium (K) were estimated by using flame photometer (Model AnA-135, OSK, Japan). Most of the elements like Calcium, Magnesium, Iron(Fe), Zinc (Zn), Copper (Cu), Nickel (Ni), Lead (Pb) and Manganese^[26] in *C. Sophera* seed oil cake were analyzed by using Atomic Absorption Spectrophotometer (Varian, AA 240FS, Australia) which was equipped with flame and graphite furnace. For the experiment, air acetylene flame mode were used. The condition fixed with acetylene 1.8 l/min and air 15 l/min, argon gas flow for inert atmosphere. The instrumental

default temperature parameters were automatically fixed for each element analysis.

2.4.7. Carbohydrate Content

Carbohydrate content of *C. Sophera* seed oil cake was estimated by subtracting the sum of the protein, fat, ash and crude fiber from the dry sample.

$$\text{Carbohydrate (\%)} = 100 - (\text{Protein} + \text{Fat} + \text{Ash} + \text{Crude Fibre}) \times 4 \text{ cal/g}$$

3. RESULTS AND DISCUSSION

Proximate composition of *C. Sophera* seed oil cake was recorded and the results are presented in Table 1 and Figure-1 respectively.

Table 1: Proximate composition (%) *C. Sophera* seed oil cake.

Test parameters	<i>C. Sophera</i> seed oil cake
	Percent (%) Composition
Moisture	11.53 ± 0.23
Organic matter	88.47 ± 0.23
Ash on drying	38.55 ± 0.10
Water soluble part	6.72 ± 0.02
Water insoluble part	87.00 ± 0.02
Carbohydrate	38.65 ± 0.30

Data are expressed as Mean ± SD (n=3).

C. Sophera seed oil cake has high ash content (384.55%), which indicates the presence of high quality of mineral contents. The moisture content for this oil cake was determined on the fresh weight basis whereas the organic content was calculated on the dry weight basis. Water soluble ash indicates the content of soluble minerals.^[24] The present study showed water soluble part for *C. Sophera* seed oil cake is 6.72% and water insoluble part is 87%.




Figure 1: Proximate analysis of *Cassia Sophera* seed oil cake.

Plants require energy in chemical form so they can grow and carry out basic life functions like all living organisms. Plants produce, store and burn carbohydrates in the form of sugar to provide themselves with

energy.^[25] The present study showed the presence of carbohydrate content in the *C. Sophera* seed oil cake is 38.65%.

A total of 10 elements i.e. Na, K, Ca, Mg, Fe, Zn, Cu, Pb, Mn and Ni were analyzed from *C. Sophera* seed oil cake by AAS which are accountable for growth of plants in agricultural sector. The result of the analysis is presented in the Table 2 & Figure-2 respectively. It may be noted each result is an average of at least three independent measurements. For the formation of secondary metabolites which are responsible for pharmacological actions these elements play a vital role.

Table 2: Mineral compositions (mg/kg) of *C. Sophera* seed oil cake.

Elements	<i>C. Sophera</i> seed oil cake (dry weight basis, mg/kg)
Na	237.74±0.027
K	332.01±0.071
Ca	22.07±0.001
Mg	78.30±0.004
Fe	1.15±0.0001
Zn	0.061±<0.001
Cu	0.081±0.01
Ni	0.062±<0.001
Mn	0.122±0.001
Pb	0.003±<0.001

Measured values are mean ± Standard Deviation (SD) of three replicate analysis. From the present study it was observed that Potassium showed higher concentration (332.01 mg/kg) than the other elemental concentration. In photosynthesis, Potassium regulates the opening and closing of stomata and therefore, regulates CO₂ uptake. Also Potassium triggers activation of enzymes and is essential for production of ATP. Potassium plays a major role in the regulation of water in plants (osmoregulation).^[26]

Although Sodium (Na⁺) is not essential for most plants but it can be beneficial to plants in many conditions particularly when Potassium (K⁺) is deficient. As such it can be regarded as “functional nutrient”. It has been observed many times that during K⁺ deficiency many (glycophytic) plants respond positively to Na⁺ fertilization for salt-tolerant (halophytic) plants, even high concentration of Na⁺ promote growth. From the present study it was found the sodium concentration is 237.74 mg/kg in *C. Sophera* seed oil cake.^[27]




Figure 2: Mineral compositions (mg/kg) of seed oil cake.

The nutrients of plants need are categorized into three different categories, main nutrients, micro nutrients and trace elements. N, P & K are considered primary nutrients. Ca along with Fe, Zn, Mn, Mg, S etc. are micro nutrients and some elements such as Molybdenum, Nickel, Selenium etc. are considered trace elements. Even though the main nutrients are very important for most plants, Calcium is even more important for some plants like tomatoes. In the form of Calcium pectate, Calcium holds the cell walls of plants together. It also activates specific plant enzymes, which sends signals to the plant cells that coordinate growth activities. From the present study it was found the Calcium concentration is 22.07 mg/kg in *C. Sophera* seed oil cake.^[28]

Magnesium is involved in many physiological and biochemical processes; it is an essential element for plant growth and development and plays a key role in plant defense mechanisms in abiotic stress solutions. The most commonly known function of Mg in plants is probably its role as the central action of the chlorophyll molecule in the light absorbing complex of chloroplasts and its contribution to photosynthetic fixation of CO₂. From the present study it was found the Magnesium concentration is 78.30 mg/kg in *C. Sophera* seed oil cake.^[29]

Zinc is plant micro nutrient which is involved in many physiological functions. Its inadequate supply will reduce crop yields. Zinc deficiencies can affect plant by stunting its growth, decreasing number of tillers,

chlorosis and smaller leaves, increasing crop maturity period, spikelet sterility and inferior quality of harvested products. From the present study it was found the Zinc concentration is 0.061 mg/kg in *C. Sophera* seed oil cake.^[30]

Iron is an essential micro nutrient for almost all living organisms because it plays critical role in metabolic processes such as DNA synthesis, respiration and photosynthesis. From the present study it was found the Iron concentration 1.15 mg/kg in *C. Sophera* seed oil cake.^[31]

Manganese (Mn) is an essential micro nutrients in most organisms. In plants, it participates in the structure of photo-synthetic proteins and enzymes. Its deficit is dangerous for chloroplast because it affects the water-splitting system of photo-system II (PS II) which provides the necessary electrons for photo-synthesis. From the present study it was found the Manganese concentration is 0.122 mg/kg in *C. Sophera* seed oil cake.^[32]

Lead (Pb) causes a number of toxicity symptoms in plants e.g. stunted growth, chlorosis and backening of root system. Pb inhibits photo-synthesis, upsets mineral nutrition and water balance. From the present study it was found the Lead concentration is 0.003 mg/kg in *C. Sophera* seed oil cake which lies in the permissible range.^[33]

Copper (Cu) is an essential metal for normal plant growth and development, although it is potentially toxic. Cu participates in numerous physiological processes and is essential co-factor for many metalloproteins, however problem arises when excess Cu is present in cells. Excess Cu inhibits plant growth and impairs important cellular processes. From the present study it was found the Copper concentration 0.081 mg/kg in *C. Sophera* seed oil cake.^[34]

4. CONCLUSION

From the above study (Proximate analysis & elemental analysis) suggested that the *C. Sophera* seed oil cake contain sufficient amount of macro and micro elements which justify its suitability in agro-industrial uses. *C. Sophera* seed oil cake can be used for the production of biofertilizer in agricultural sectors. However more detailed analysis of chemical composition of *C. Sophera* seed oil cake is required.

ACKNOWLEDGEMENT

We are grateful to INARS, BCSIR for giving us the opportunity to do mineral compositions analysis by AAS of plant materials. We are also thankful to the Director, BCSIR Laboratories, Dhaka for providing necessary facilities to carry out this research work.

REFERENCES

1. Ayeni, MJ, Obeyemi, SD, Kayode, J, Peter, GP. Phytochemical, Proximate and Mineral Analysis of the leaves of *Gossypium hirsutum* L., and

- Momordica charantia* L., *Journal of Natural Science and Research*, 2015; 6(5).
2. Drever, BD, Anderson, WG, Riedel, G, Kim, DH, Ryu, JH, Choi, DY, Platt, B. The seed extract of *Cassia Obtusifolia* offers neuroprotection to mouse hippocampal cultures, *J. Pharmacol Sci.*, 2008; 107: 380-92.
3. Guo H, Chang Z, Yang R, Guo Dm Zheng J. Anthraquinones from hairy root cultures of *Cassia Obtusifolia*, *Phytochemistry*, 1998; 49: 1623-1625.
4. Aminabee, SK and Lakshmana R. A Plant Review of *Cassia Sophera* Linn. *International Journal of Pharmaceutical, Chemical and Biological Sciences*, 2012; 2(3): 408-414.
5. Boraste, A, Vamsi, KK, Jhadav, A, Khairnar, Y, Gupta, N, Trivedi, S, Patil, P, Gupta, G, Gupta, M, Mujapara, AK and Joshi, B. Biofertilizers: A novel tool for agriculture. *International J. f Microbiol. Res.*, 2009; 1(2): 23-31.
6. Abdou, MA and Mahmoud, AH. Growth and oil production of *Foeniculum vulgare* Mill. 2. The effect of number of irrigation and organic fertilizers. *J. Agric. Sci. Mansoura, Univ.*, 2003; 28(5): 3868-3888.
7. Hassan, FAS and Ali, EF. A comparative study between traditional mineral nutrition and other alternative sources on anise plant. *Europe. J. Sci. Res.*, 2013; 106(2): 201-212.
8. Fernandes, DM, Doi, NT, Villas, RL, Bull, LT, SS and Miazaki. Macronutrients and micronutrients removal by aerial part of sweet fennel due to liming and chicken manure fertilization. *Acta Hort.*, 2002; 569: 185-190.
9. Yasin, M, Ahmad, K, Mussarat, W and Tanveer, A. Bio-fertilizers, substitution of synthetic fertilizers in cereals for leveraging agriculture. *Crop Environ.*, 2012; 3: 62-66.
10. Gendy, ASH, Said-Al Ahl, HAH and Mahmoud, AA. Growth, Productivity and Chemical Constituents of Roselle (*Hibiscus sabdariffa* L.) Plants as Influenced by Cattle Manure and Biofertilizers Treatments. *Australian. J. Basic. Appl. Sci.*, 2012; 6(5): 1-12.
11. Hassan, FAS, Ali, EF and Mahfouz, SA. Comparison between different fertilization sources, irrigation frequency and their combinations on the growth and yield of coriander plant. *Australian J. Basic. Appl. Sci.*, 2012; 6(3): 600-615.
12. Chen, J. The combined use of chemical and organic fertilizers and/or biofertilizer for crop growth and soil fertility. International Workshop on Sustained Management of the Soil Rhizosphere System for Efficient Crop Production and Fertilizer Use., 2012; 16-20 October, Thailand.
13. Hellal, FA, Mahfouz, SA and Hassan, FAS. Partial substitution of mineral nitrogen fertilizer by biofertilizer on (*Anethum graveolens* L.) plant. *Agri. Biol. J. North America*, 2011; 2(4): 652-660.
14. Akhani, A, Darzi, MT and Hadi, MHS. Effects Of Biofertilizer And Plant Density On Yield

- Components And Seed Yield Of Coriander (*Coriandrum Sativum*). *Inter. J. Agricult. Crop Sci.*, 2012; 4-16: 1205-1211.
15. Sokhangoy, SH, Ansari, KD and Asli, E. 'Effect of bio-fertilizers on performance of dill (*Anethum graveolens*). *Iranian. J. Plant Physiol.*, 2012; 2(4): 547-552.
 16. Dadkhah, A. Effect of Chemicals and Bio-fertilizers on Yield, Growth Parameters and Essential Oil Contents of Fennel (*Foeniculum vulgare* Miller.). *Journal of Medicinal Plants and By-products.*, 2012; 2: 101-105.
 17. Al-Fraihat, AH, Al-dalain, SYA, AlRawashdeh, ZB, Abu-Darwish, MS and Al-Tabbal, JA. Effect of organic and biofertilizers on growth, herb yield and volatile oil of marjoram plant grown in Ajlloreion, *Jordan. J. Medicinal Plants Res.*, 2011; 5(13): 2822-2833.
 18. Shaalan, MN. Influence of biofertilization and chicken manure on growth, yield and seeds quality of *Nigella sativa*, L. plants. *Egypt, J. Agric. Res.*, 2005; 83(2): 811-828.
 19. Valadabadi, SA and Farahani, HA. Influence of biofertilizer on essential oil, harvest index and productivity effort of black cumin (*Nigella sativa* L.). *Inter. J. Biotechnol. Mole. Biol. Res.*, 2013; 4(2): 2427.
 20. United States of Pharmacopeia United States of Pharmacopeia national Formulary USP30-NF25. Zgeneral test and Assay, USP method 56.1.12601 Twin Brook parkway Rockville, Maryland 20852-1790, USA. 2007.
 21. Osborn, DR, Voogt, P. Calculation of Calorific value., In: *The Analysis of Nutrients in Foods*; Academic press, New York, 1978; pp. 239-240.
 22. Anjana, B and Metai, S. Composition of Indian Aquatic Plants in relation to utilization as Animal Forage. *J. Aquat. Plant Manage*; 1990; 28: 69-73.
 23. The Wealth of India. First supplement series (Raw materials). 2004; Vol.1: A-Ci, NISCAIR, CSIR, New Delhi, 7236-208.
 24. Kerry, HC, Debra, M. The Structure, function and biosynthesis of plant cell wall pecticpolysaccharides. *Carbohydrate Research*, 2009; 344: 1879-1900.
 25. Vander, AJ, Luciano, D, Sherman J.H. Human physiology: The Mechanism of body function, 2001; 8th ed. Boston: Mc-Graw Hill.
 26. Kalavati, P and Modi, HA. The Importance of Potassium in Plant growth-A review, *Indian Journal of Plant Sciences*, 2012; 1(02-03): 177-186.
 27. Matthius, FJM. Sodium in plants, perception, signaling and regulation of sodium fluxes, *J. of Experimental Botany*, 2014; 65(3): 849-858.
 28. Philip, JW and Martin, R B. Calcium in Plants, *Annals of Botany*, 2003; 92: 487-511.
 29. Mehmet, S, Andreas, G, Verena, W and Heike, T. Role of Magnesium fertilizers in agriculture Plant soil continuum, *Crop and Pasture Sciences*, 2015; 66: 1219-1229.
 30. Hafeez, B, Khanif, YM and Saleem, M. Role of Zinc in Plant Nutrition- A review, *American J of Experimental agriculture*, 2013; 3(2): 374-391.
 31. Gyana, RR and Sumita, S. Role of iron in plant growth and metabolism, *Reviews in Agricultural Science*, 2005; 3: 1-24.
 32. Milaleo, R, Reyes-Diaz, M, Ivanov, AG, Mora, MI and Alberdi, M. Manganese an essential and toxic element for plants: Transport, accumulation and resistance mechanisms, *J. Soil Sci. Plant Nutr.*, 2010; 10(4): 476-494.
 33. Pallavi, S and Ruma, SD. Lead toxicity in plants, *Braz. J. Plant Physiol*, 2005; 17(1): 35-52.
 34. Inmaculada Y. Copper in Plants, *Braz. J. Plant Physiol*, 2005; 17(1): 145-156.